



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

The role of biomarkers in diagnosis of COVID-19 – A systematic review

Muhammed Kermali, Raveena Kaur Khalsa, Kiran Pillai, Zahra Ismail, Amer Harky



PII: S0024-3205(20)30536-1

DOI: <https://doi.org/10.1016/j.lfs.2020.117788>

Reference: LFS 117788

To appear in: *Life Sciences*

Received date: 8 May 2020

Accepted date: 11 May 2020

Please cite this article as: M. Kermali, R.K. Khalsa, K. Pillai, et al., The role of biomarkers in diagnosis of COVID-19 – A systematic review, *Life Sciences* (2018), <https://doi.org/10.1016/j.lfs.2020.117788>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The role of Biomarkers in diagnosis of COVID-19 – A systematic review

Muhammed Kermali¹, Raveena Kaur Khalsa¹, Kiran Pillai¹, Zahra Ismail¹ Amer Harky²

1. St. George's, University of London, Cranmer Terrace, London SW17 0RE
2. Department of Cardio-thoracic Surgery, Liverpool Heart and Chest Hospital, Liverpool, UK

Corresponding author:

Amer Harky

MRCS, MSc

Department of Cardiothoracic Surgery

Liverpool Heart and Chest Hospital

Liverpool, UK

E-mail: aaharky@gmail.com

Tel/fax: +44-151-600-1616

Conflict of Interest: none to be declared

Funding: No funding obtained

Word Count: 3,416

Word Count of Abstract: 197

Number of Tables and Figures: 11

Key words: SARS-CoV-2, COVID-19, Biomarkers, Blood tests

Abstract**Aims**

As of the 28th April 2020, the COVID-19 pandemic has infiltrated over 200 countries and affected over three million confirmed people. We review different biomarkers to evaluate if they are able to predict clinical outcomes and correlate with the severity of COVID-19 disease.

Methods

A systematic review of the literature was carried out to identify relevant articles using six different databases. Keywords to refine the search included 'COVID-19', 'SARS-CoV2', 'Biomarkers', among others. Only studies which reported data on pre-defined outcomes were included.

Key Findings

Thirty-four relevant articles were identified which reviewed the following biomarkers: C-reactive protein, serum amyloid A, interleukin-6, lactate dehydrogenase, neutrophil-to-lymphocyte ratio, D-dimer, cardiac troponin, renal biomarkers, lymphocytes and platelet count. Of these, all but two, showed significantly higher levels in patients with severe complications of COVID-19 infection compared to their non-severe counterparts. Lymphocytes and platelet count showed significantly lower levels in severe patients compared to non-severe patients.

Significance

Although research is still in its early stages, the discovery of how different biomarkers behave during the course of the disease could help clinicians in identifying severe disease earlier and subsequently improve prognosis. Nevertheless, we urge for more research across the globe to corroborate these findings.

Introduction

On December 2019, Wuhan City in China, became the epicentre of unexplained cases of pneumonia. On January 2020, Chinese scientists identified this as a novel coronavirus, temporarily labelled as, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)(1). Its name was then changed to coronavirus disease 2019 (COVID-19) by the World Health Organization in February 2020 as the disease spread worldwide (2).

It has been suggested the outbreak has a zoonotic origin, and like other respiratory pathogens, it spreads through human-to-human transmission such as coughing and sneezing(3). Although limited, research suggests a possibility of transmission even amongst the asymptomatic(4).

As of the 28th April 2020, over 200 countries have been affected by the COVID-19 disease with over three million confirmed cases leading to over 200,000 deaths. Yet it is believed many remain unreported in certain areas of the world(1).

Infected patients may present with any of the following; fever, high temperature (>37.3 C), cough, myalgia, sputum production, headache, haemoptysis, diarrhoea, dyspnoea and in some cases, acute respiratory distress syndrome (ARDS), acute cardiac injury or secondary infection(5).

Upon examination, these subjective clinical symptoms can be interpreted more confidently with the use of biological markers (biomarkers). These provide objective values throughout the progression of the disease(6). Henceforth, categorising patients into mild, severe or critical becomes more defined, allowing for earlier interventions (7).

This article aims to explore the role of different biomarkers in the disease pathogenesis of COVID-19 and assess how their levels vary depending on the severity of the disease. By doing so, it gives clinicians a tool to group patients and predict prognosis and mortality. The biomarkers we review include, C-reactive protein (CRP), IL-6, white cell count (WCC), lactate dehydrogenase (LDH), D-dimers, platelet count, cardiac troponin and renal markers.

Methods and Materials

1. Search Strategy

A comprehensive literature search was done on PubMed, SCOPUS, Embase, Cochrane database, Google Scholar and Ovid to identify articles discussing biomarkers in this review and its clinical implications on COVID-19 in accordance with Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines. Key words used were 'C-reactive protein', 'COVID-19', 'interleukin-6', 'lactate dehydrogenase', 'SARS', 'white cell count', 'neutrophil count', 'lymphocyte count', 'D-dimer', 'platelet count', 'cardiac troponin', 'renal biomarkers', 'urea', 'creatinine'. The search terms were used as key words and in combination as MeSH terms to maximize the output from literature findings. A staged literature search was done, whereby a separate literature search was performed for each section within this article and all the relevant studies were identified and summarized separately. If a paper reports on multiple biomarker, then the results have been shared between different parts of this review. The relevant articles are cited and referenced within each section separately. No limit has been placed on publication time or language of the article. All the relevant articles were identified and screened by three authors; the results are summarized in narrative manner in each relevant section of this review. A summary table of each section is provided where appropriate.

2. Inclusion and exclusion criteria:

Studies were included if they have reviewed a correlation between a biomarker and the severity of COVID-19. Exclusion criteria were editorials, consensus documents, commentaries, and studies with no particular definition of the role of biomarkers in COVID-19.

3. Data extraction:

All articles were screened by two authors and any disagreement was reached by consensus or involvement of third author. Data extracted by two authors and validated by third author.

4. Quality assessment:

Quality of each publication was evaluated by two independent reviewers. This review addressed key domains: type of biomarker, level of the biomarker, correlation with severity of the disease, survival, number of patients investigated and outcomes.

5. Statistical analysis:

It was not possible to conduct an appropriate meta-analysis because there were not enough research data among the studies on this subject.

Results

A total of 414 articles were found. After removal of duplicates, 224 articles were used for full-text screening with 34 studies included in our analysis. The complete PRISMA flow chart is reported in figure 1. Biomarkers measured were C-reactive protein (CRP), interleukin-6 (IL-6), White cell count (WCC), lactate dehydrogenase (LDH),

D-dimer, platelet count, high-sensitivity troponin I (hs-TnI) and renal markers. Table 1 summarises the study's characteristics.

C-reactive protein

CRP is a plasma protein produced by the liver and induced by various inflammatory mediators such as IL-6. Despite being non-specific, this acute phase reactant is used clinically as a biomarker for various inflammatory conditions; a rise in CRP levels are associated with an increase in disease severity(7).

The application of CRP in COVID-19 has been highlighted by a retrospective single-centre study in Wuhan, China, where the majority of patients in the severe cohort showed significantly higher levels compared to the non-severe cohort (57.9 mg/L vs 33.2 mg/L, $P < 0.001$)(8). A second retrospective cohort study found the likelihood of progressing to severe COVID-19 disease increased in patients with CRP levels greater than 41.8mg/L(9). Both studies suggest CRP levels are a strong indicator to reflect the presence and severity of COVID-19 infection.

Furthermore, a study from unpublished observations suggests CRP is one of the first biomarkers within blood plasma that changes to reflect physiological complications; if accepted CRP will be the most effective biomarker to predict the progression of COVID-19 infection. Contrastingly, the same study illustrated some cases of infection which showed changes in serum amyloid A (SAA) instead of evoking significant CRP changes thus requiring further evaluation (10).

Whilst the use of SAA as a biomarker for COVID-19 requires further research, CRP and SAA are commonly used in conjunction to monitor inflammatory diseases. Though SAA is another acute phase reactant, it is responsive to both viral and bacterial infections compared to CRP(11).

Pathologically, computed tomography (CT) scans can identify lung lesions relating to COVID-19. Nonetheless, a study conducted in China revealed CT scores could not differentiate mild cases from severe. However, compared to erythrocyte sedimentation rate (ESR), CRP levels were significantly greater during early periods of severe cases and proved to be a more sensitive biomarker in reflecting disease development(12).

The excellent performance of CRP as a biomarker is reflected in the 'area under curve' in the receiver operating analysis of 0.87 (95% CI, 0.10–1.00) where values 83% and 91% represent sensitivity and specificity, respectively. Hence compared to CT scans alone, CRP values are more reliable for earlier identification of case severity (12). Table 2 summarises the studies used in analysing CRP and COVID-19.

Interleukin-6

Cytokine release syndrome (CRS) is an over-exaggerated immune response involving an overwhelming release of pro-inflammatory mediators. This mechanism underlies several pathological processes including acute respiratory distress syndrome (ARDS) (13). Studies investigating the role of cytokines in SARS and MERS have also found a link between CRS and disease severity(14). Understanding their role in COVID-19 disease may help facilitate the design of novel immunotherapies.

Studies have revealed that levels of IL-6, the most common type of cytokine released by activated macrophages, rise sharply in severe manifestations of COVID-19(15). However, since most studies to date have been observational, it is difficult to

extrapolate if the rise is significant enough to cause the manifestations seen in severe forms.

One meta-analysis reviewing six studies show mean IL-6 concentrations were 2.9-fold higher in patients with complicated COVID-19 compared to those with non-complicated disease (n=1302; 95% CI 1.17-7.19)(16). In its analysis, the outcomes of the studies include ICU admission, onset of ARDS and mortality. Since the proportionate rise of IL-6 is correlated with disease severity, this study can prove ground-breaking. Although clinicians can use this to identify severity earlier and commence oxygen therapy sooner, the varying outcomes makes it somewhat difficult to ascertain what level of IL-6 corresponds to what negative outcome. Furthermore, many studies recruited participants from the same centre, giving rise to the potential of selection bias. Table 3 summarises the studies used in analysing IL-6 and COVID-19.

White Cell Count

White blood cells (WBCs), known as leucocytes are a component of blood generated from bone marrow and lymphoid tissue. They are divided into two major groups, granulocytes and agranulocytes. Within the granulocyte group are eosinophils, basophils and neutrophils (NC), whereas lymphocytes (LC) and monocytes are present in agranulocytes. A disproportionate number of these cells may reveal an underlying infection and hence can be measured using blood tests, producing a WCC. However, the reliability of WCC as a biomarker for COVID-19 remains unproven.

A retrospective study found several differences in WCC between severe and non-severe COVID-19 patients(8). Both groups experienced an increase in leucocytes with the severe group having a significantly greater rise (5.6 vs $4.9 \times 10^9/L$; $P <$

0.001). NCs were predominantly driving this increase as the severe set (4.3 vs $3.2 \times 10^9/L$; $P < 0.001$). Interestingly, the levels of lymphocytes, monocytes, basophils and eosinophils were less, resulting in a greater neutrophil-to-lymphocyte ratio (NLR; 5.5 vs 3.2 ; $P < 0.001$). NLR is an infamous biomarker, high in wide-spread inflammatory conditions and can be used to reflect disease severity. However, a larger study is needed to clarify NLR's effectiveness as a biomarker.

Another conducted in China concludes similar findings of high NC and low LC count in severely affected patients, suggesting NLR could be a potential biomarker for early detection of severe COVID-19 (17).

However, other factors may disrupt the accuracy of the WCC results observed. These include glucocorticoid therapy and other underlying viral/bacterial infections(11).

LC is separately addressed in the literature, secondary to NLR. A descriptive study in China reported depleted LC levels in the majority of COVID-19 patients(15).

Another study has found low blood lymphocyte percentage (LYM%) in critically ill patients, suggesting low LC count indicates poor prognosis. However, since the virus can target lymphoid tissue and mechanisms of IL-6, other causes of low LC count must be investigated(18). Similar to NLR, the clinical benefits of LC count as a biomarker for COVID-19 remains uncertain.

Table 4 summarises the studies used in analysing WCC and COVID-19.

Lactate Dehydrogenase

In glucose metabolism, the enzyme LDH converts pyruvate to lactate. LDH secretion is triggered by necrosis of the cell membrane, hinting to viral infection or lung

damage, such as the pneumonia induced by SARS-CoV-2 (19). There is convincing evidence linking LDH levels to the development of COVID-19 disease (20).

A study found significantly higher levels of LDH in ICU patients than non-ICU patients (248 U/L vs 151 U/L, $p=0.002$). Since high levels of LDH continued in the ICU patients number of days post-admission (160 U/L vs 218 U/L, $p=0.002$), LDH may be a predictive biomarker of severe disease. However, the one centre study may be prone to selection bias which could potentially reduce its validity (21).

A multi-centre study involving 1099 patients reported supporting evidence correlating extent of tissue damage and inflammation with increasing levels of LDH(22). Furthermore, when LDH levels were correlated with CT scans, significantly higher levels reflected the severity of pneumonia(23).

There is increasing confidence in using LDH as a biomarker to measure severity of COVID-19 infection. Another study found that there was a significant rise in LDH levels among refractory COVID-19 patients (24).

Table 5 summarises the studies used in analysing LDH and COVID-19.

D-dimer

D-dimer originate from the lysis of cross-linked fibrin with rising levels indicating the activation of coagulation and fibrinolysis(25). Early studies have associated COVID-19 with haemostatic abnormalities with one study observing elevated levels of D-dimer, the measure of coagulation, in non-survivors compared to survivors(26).

A retrospective cohort study composed of 191 patients found that D-dimer levels greater than 1.0 $\mu\text{g/ml}$ ($p=0.0033$) were associated with increased mortality among COVID-19 patients. Furthermore, they found that levels of 2.0 $\mu\text{g/ml}$ or more on admission was the optimum cut-off to predict in-hospital mortality for COVID-19(27).

Studies have reported that nearly 90% of inpatients with pneumonia had increased coagulation activity marked rising D-dimer levels(28).

Furthermore, Huang et al. found that levels of D-dimer on admission could be used to triage patients into critical care (5). The researchers found that median D-dimer levels were higher in ICU patients compared to non-ICU patients (2.4 mg/L vs. 0.5mg/L; $p=0.0042$). This, along with the previous study, suggest that D-dimer levels can be used as a prognostic marker and help clinicians monitor those who are likely to deteriorate earlier. However, this study confirmed the diagnosis of COVID-19 using lower respiratory tract specimens and did not use paired nasopharyngeal swabs to investigate the viral RNA detection rate between the upper and lower respiratory tract specimens. Secondly, with a cohort size of 41 patients, it is difficult to assess predictors of disease severity and mortality with multivariable-adjusted methods.

Table 6 summarises the studies used to analyse D-dimer and COVID-19.

Platelet count

As seen with previous coronavirus outbreaks, COVID-19 infection leads to severe haematological changes leading to thrombocytopenia.

Meta-analysis of 1799 patients reveal those with severe COVID-19 infections had significantly lower platelet counts (WMD $-31 \times 10^9/L$; 95% CI, -35 to $-29 \times 10^9/L$)(29). When using mortality as an endpoint, non-survivors evidently had a significantly lower platelet count (WMD, $-48 \times 10^9/L$; 95% CI, -57 to $-39 \times 10^9/L$). Using thrombocytopenia as an endpoint also revealed a fivefold greater risk of COVID-19 (OR, 5.13; 95% CI, 1.81-14.58). Despite the varying definitions of disease

severity and thrombocytopenia having an influence on results analysis, platelet count could possibly be used clinically to indicate infection severity.

A retrospective study which used Cox proportional hazard regression analysis found that platelet count is an independent risk factor for mortality among COVID-19 patients, where a $50 \times 10^9/L$ increase is associated with 40% decreased mortality (HR 0.60, 95%CI: 0.43, 0.84)(30). Here, thrombocytopenia at admission was more likely to occur in non-survivors than in survivors. Although many risk factors were accounted for in this study, the possibility for unmeasured confounder cannot be excluded.

Another study corroborates the previously documented work. The nadir platelet count was significantly associated with mortality – and the lower the nadir, the stronger the association(31). Again, thrombocytopenia was more likely to occur in non-survivors than survivors. This study is from adequate sample sizes providing statistical power, however, similar to the previous studies, they are all retrospective making the correlation seen difficult to extrapolate from.

Testing the platelet count is a routine part of laboratory tests and the literature suggests it has inherent value in providing more detail on the patient's condition.

Table 7 summarises the studies used in analysing platelet count and COVID-19.

Cardiac Troponin

There is growing evidence of higher mortality rates among those with underlying cardiovascular disease due to COVID-19 infection (22). Some have investigated the use of high-sensitivity cardiac troponin I (hs-TnI) as a marker of disease progression and mortality.

A retrospective study performed in China of patients with confirmed COVID-19 based on SARS-CoV-2 RNA detection, revealed a univariable odds ratio for death at 80.1 (95% CI 10.3-620.4, $p < 0.0001$) for hs-TnI(32). This risk was higher compared to other biomarkers such as D-dimer and lymphocyte count. Another study of 416 hospitalised patients with COVID-19 reported that hs-TnI was elevated in 1 in 5 patients on presentation(33). These patients were more likely to require invasive (22% vs 4%, $p < 0.001$) or non-invasive (46% vs 4% , $p < 0.001$) ventilation, develop ARDS (59% vs 15%, $p < 0.001$) or acute kidney injury (9% vs 0%, $p < 0.001$).

Early recognition of myocardial injury indicated by elevated hs-TnI aid in appropriate triage to a critical care area and inform the use of inotropes and vasopressors. However, elevated levels are common in hospitalized patients and are likely to be due to non-ischaemic causes of myocardial injury. This may lead to inappropriate use of cardiology consultation and downstream testing and increased risk to cardiac physiology staff.

Table 8 summarises the studies used in analysing hs-TnI and COVID-19.

Renal markers

There is also evidence that chronic kidney disease is associated with severe forms of COVID-19 infection(34).

Studies have demonstrated significantly higher levels of renal biomarkers such as serum urea, creatinine and markers of glomerular filtration rate in severe cases(35). Since these results stem from the analysis of 28 patients, extrapolation across larger cohorts is more difficult.

A larger study of 701 patients revealed that elevated serum creatinine levels on admission correlated with severity due to significant abnormalities in the coagulation

pathway(36). They also found that these patients were more likely to require mechanical ventilation or be placed in intensive care. Univariate Cox regression analysis found elevated creatinine levels was also associated with in-hospital mortality (HR 2.99, 95% CI: 2.00, 4.47). Proteinuria, haematuria and elevated urea levels had similar, if not larger, hazard ratios.

Interestingly, another study showed a potential role for urinalysis over serum markers of kidney function(37). Here, abnormalities in the routine urine test on admission correlated strongly with disease severity. They go on to suggest that urinalysis may reveal kidney impairment more readily than evaluation of serum renal biomarkers. However, these tests were only carried out on admission and so patients in earlier stages of the infection had changes in serum levels obscured by compensatory kidney function. Hence renal abnormalities on admission may indicate higher risks of deterioration, ensuring appropriate triaging.

Table 9 summarises the studies used in analysing renal markers and COVID-19.

Discussion

COVID-19 is a rapidly spreading pandemic increasing the burden on medical facilities. Symptoms vary from mild fever to ARDS complicating diagnosis, prognosis, and monitoring. Hence it is vital to ascertain a patient's condition in a timely manner. Biomarkers are quantitative measurements used clinically for many conditions reflecting pathological development. A summary of the biomarkers discussed in this review can be found in Table 10.

When assessing a patient with COVID-19 infection, biomarkers can be useful to clinicians in starting treatment and close monitoring. Though biomarkers may help

improve prognosis and outcomes, their significant variability between patients could affect the findings of the studies.

Compared to other biomarkers, there is hesitancy in using WCC alone as it is influenced by many factors such as glucocorticoid treatment which increases it. Although WCC encompasses many cell types, NCs and LCs are most clinically relevant biomarkers. Multiple studies on COVID-19 have concurred high NLR in severe cases compared to non-severe cases due to high NC and low LC. The use of LC independently has been suggested as a potential biomarker of COVID-19 as patients have consistently low LC, with significant lymphopenia reported in critically ill patients(18). Therefore, further research on WCC accuracy for assessing disease progression is necessary.

Although, most of the studies referenced in this review are single centred studies originating in Wuhan, China, the virus is now a global pandemic thus requiring international studies. As a copious amount of data was collected from the same location, it is possible patients may have been used in more than one study. Furthermore, categorising patients into severe and non-severe sub-groups is dependent on the definitions of severity of COVID-19. Two very different measures of severity used in some studies are ICU admission and onset of ARDS. This questions the heterogeneity across the studies and validity of the findings.

Variability can be accounted to the differences in patient exposure to the virus or intraindividual variability, where differences in laboratory procedures lead to errors. Consequently, the sensitivity and specificity of biomarkers are useful indicators to the effectiveness, though not all studies discussed this. To ascertain the usefulness of the biomarkers listed in this review as indicators of disease progression, and whether

they definitively rise in COVID-19 requires further data collection. Statistical analysis using receiver-operator characteristic curves can also provide analysis sensitivity and false-positive rates.

Findings of these studies has discovered changes in biomarker levels and may potentially be useful in creating a therapeutic intervention. For instance, one study has reviewed the use of anticoagulation therapy in patients with coagulopathy or marked rise in D-dimers in the setting of COVID-19(38). Low molecular weight heparin was found to be associated with better prognosis in severe cases. To further asses the role of anti-coagulants as a treatment, we encourage large interventional trails to study this.

Many studies in this review were limited due to small sample size and selection bias if conducted at one centre. To improve reliability and reproducibility, more research into the prognostic value of biomarkers is necessary.

Conclusion

In conclusion, the work to date suggests that there is clear evidence of how the levels of biomarkers may change according to severity of COVID-19 infection. This can be used as an adjunct in clinical practice to guide treatment and admission to ICU. By doing so, it may improve prognosis and minimise the mortality rates. However, being in the infant stages of understanding the pathology of this infectious disease, we urge for further research worldwide to better understand the changes noted in this review.

Conflict of Interest Statement

The authors declare that there are no conflicts of interest.

Table 1 – Study Characteristics

Author	Study design	Country	Cohort size	Biomarkers studied	Comments
Huang et al. (2020) (5)	Prospective	China	41	Lymphocytes, IL-6, D-dimer, platelet count	Compared to non-ICU patients, ICU patients had higher plasma levels of IL-6, D-dimer and platelets. Lymphopenia was more common in ICU patients than non-ICU patients
Li et al. 2020 (39)	Retrospective cohort; single centre	China	132	CRP, WCC	Critically severe patients had significantly higher CRP and WCC than severe or non-severe patients.
Liu et al (9)	Retrospective cohort; single centre	China	140	CRP, IL-6	Significantly more patients in the severe group experienced higher CRP and IL-6 levels vs non-severe.
Ji et al. 2020 (10)	Retrospective cohort; single centre	China	33	CRP, D-dimer, LC	The preprint study shows CRP, D-dimer significantly increased but LC decreased for patients in direct vs indirect contact to Wuhan – does not assess severity. CRP levels compared by ratio to other blood parameters indicated CRP changes prior to others.
Tan et al. 2020 (12)	Retrospective cohort; single centre	China	27	CRP, WCC	ROC analysis showed strong association found between CRP levels and progression of disease. Interestingly, absolute values of increased WCC and

					NLR found in influenza A or B group vs SARS-COV-2.
Wang et al. 2020 (40)	Retrospective cohort; single centre	China	27	CRP	Greater CRP values correspond with the critical group, as groups were determined based on the diameter of largest lung lesion - CRP levels may indicate lung damage and development of disease.
Qin et al. 2020 (8)	Retrospective cohort; single centre	China	452	CRP, IL-6 and LDH, WCC	Significantly higher levels of CRP, IL-6, LDH and NC but low LC in severe COVID-19,vs non-severe group. Surveillance may help in early screening of critical illness
Wang et al. 2020 (17)	Retrospective cohort; single centre	China	138	WCC, NC, LC, LDH, D-dimer, blood urea, creatinine	Higher WCC, specifically higher NC, lower LC and higher NLR in ICU vs non-ICU patients. Prior to the death of critical patients: NC, D-dimer, blood urea, and creatinine levels rose throughout until death, whilst the LC carried on falling. Elevated LDH also correlated with patients in ICU vs non-ICU.
Chen N et al. 2020 (15)	Retrospective cohort; single centre	China	99	WCC, NC, LC, LDH, CRP, platelet count, D-dimer, IL-6, blood urea, serum creatinine	Levels of LDH, CRP,D-dimer, IL-6 increased over normal ranges in COVID-19- induced pneumonia patients, whereas increases and decreases were

					seen in WCC (high NC, low LC), blood urea, serum creatinine and platelet count. Absolute LC values decreased across patients; destruction of T lymphocytes may lead to worse outcomes.
Tan et al. 2020 (18)	Retrospective cohort; single centre	China	90	WCC	Graphical representation of time-lymphocyte% model (TLYM%) using 5 randomly selected individuals to illustrate disease progression until death. TLYM % used for 90 hospitalised patients, it was able to categorise patients consistently. Surveillance may identify critically ill patients earlier.
Gong et al. 2020 (7) preprint	Retrospective cohort; single centre	China	100	WCC, CRP	Preprint corresponds with other studies that a high WBC, NC and CRP suggests greater severity of disease.
Luo et al. 2020 (21)	Retrospective cohort; single centre	China	35	LDH	Increased LDH levels are associated with patients in the severe group, indicating LDH may be reflective of severity of disease.
Xiong et al. 2020 (23)	Retrospective cohort; single centre	China	42	LDH	Correlates LDH values with CT scores but no breakdown is present
Mo P et al. 2020 (24)	Retrospective; single centre	China	155	LDH, NC, CRP, platelet count	LDH, NC, CRP and platelet count were higher in refractory vs general patients. Refractory

					paitents had more cases of lung abnormalities, suggesting these biomarkers correlate with development of disease
Ferrari D et al. (2020) (20)	Retrospective; single centre	Italy	141	LDH	LDH measured in COVID-19 positive vs negative patients and higher levels apparent in positive groups.
Tang et al. 2020 (26)	Retrospective cohort; single centre	China	183	D-dimer	Abnormal coagulation results with markedly elevated D-dimer are common in deaths with COVID-19
Zhou et al. 2020 (32)	Retrospective cohort; multi-centre	China	191	D-dimer, hs-Tnl, platelet count	D-dimer levels greater than 1µg/mL can help clinicians in identifying patients with poor prognosis at earlier stage There were significantly higher levels of platelets and hs-Tnl in non-survivors compared to survivors
Guan et al. 2020 (22)	Retrospective cohort; single centre	China	1099	D-dimer, Platelet count, CRP lymphocytes and LDH	Although there were evident differences in lymphocytes (lymphocytopenia), platelet count (rose) and D-dimer (rose) in patients who experienced composite endpoints (ICU admission, invasive mechanical ventilation and death) there was no statistical analysis performed LDH measured in 675. Higher

					LDH levels present in majority of severe patients vs non-severe.
Liu et al. 2020 (41)	Retrospective cohort; single centre	China	12	Lymphocytes, Albumin, Neutrophils, CRP, PaO ₂ /FiO ₂ , platelet count, creatinine and LDH	C _t value of virus correlated strongly with CRP, albumin and LDH among others. Murray score for ARDS correlated with the same markers.
Liu et al. 2020 (42)	Retrospective cohort; single centre	China	78	CRP, Albumin, Platelet count, D-dimer, AST, ALT, Creatinine	Increased CRP and decreased albumin strongly correlated with disease progression
Ruan et al. 2020 (43)	Retrospective cohort; single centre	China	150	WBC counts, lymphocytes, platelets, albumin, total bilirubin, urea, creatinine, myoglobin, cardiac troponin, CRP and IL-6	Cardiac troponin, Myoglobin, CRP and IL-6 significantly increased in cases with mortality
Yang et al. 2020 (44)	Retrospective cohort; single centre	China	52	Haemoglobin, lymphocytes, Platelet count, prothrombin time, bilirubin, creatinine and lactate.	Higher levels of platelets in severe group but lymphocytopenia seen as the most common marker of infection.
Young et al. 2020 (45)	Descriptive	Singapore	18	Haemoglobin, lymphocytes, platelet count, neutrophils, CRP and LDH	Lower platelets found in severe group, but statistical analysis was not performed
Liu et al. 2020 (30)	Retrospective cohort; single centre	China	383	Platelet parameters: platelet count, volume, distribution width and larger cell ratio	Thrombocytopenia associated with mortality Patients split on basis of thrombocytopenia not severity
Yang et al 2020	Retrospective	China	1476	Platelet count	Lowest platelet count associated

(31)	cohort; single centre				with mortality
Shi et al 2020 (33)	Retrospective cohort; single centre	China	416	Cardiac troponin, platelet count, creatinine kinase	Greater proportion of cardiac injury, marked as a rise in creatinine kinase and cardiac troponin, is more likely to require non-invasive/invasive ventilation compared to those without
Xiang et al. 2020 (35)	Retrospective cohort; single centre	China	28	AST, GGT, ALP, albumin, creatinine kinase, CKMB, CRP, creatinine, urea and cystatin C	Serum values of urea, creatinine and cystatin-C significantly increased in severe COVID-19.
Cheng et al. 2020 (36)	Prospective cohort; single centre	China	701	Creatinine	Raised creatinine levels associated with poor outcome in COVID infection
Zhou et al. 2020 (37)	Retrospective cohort; single centre	China	178	Urinalysis: proteinuria, haematuria, leucocyturia	Urinalysis on admission can effectively highlight kidney impairment
Chen L et al (2020) (46)	Prospective cohort; single centre	China	29	IL-6	Increased expression of IL-2R and IL-6 in serum is expected to predict the severity of COVID-19
Liu et al. (2020) (47)	Retrospective cohort; single centre	China	80	IL-6, CRP, LDH, ferritin and D-dimer	Baseline IL-6, CRP, LDH and ferritin was closely related to severity of COVID-19. Elevated IL-6 was significantly related to clinical manifestations of severe type patients
Diao et al (2020) (48)	Retrospective cohort; multi-	China	552 COVID; 40 healthy	T-cells, IL-6	T cells are reduced significantly and negatively correlated to IL-6

	centre				in COVID-19
Wu et al (2020) (49)	Retrospective cohort; multi-centre	China	150	IL-6, D-dimer, LDH, neutrophil count	ARDS development in COVID-19 is significantly correlated with rise in IL-6, D-dimer, LDH and neutrophil count
Zhang et al (2020) (50)	Retrospective; single centre	China	343	D-dimer	D-dimer on admission of greater than 2.0 µg/mL could effectively predict in-hospital mortality in patients with COVID-19 and could be an early and helpful marker to improve management

IL-6 = interleukin-6; ICU = intensive care unit; CRP = C-reactive protein; WCC = white cell count; LC = leucocyte count; ROC = Receiver operating characteristic; NLR = neutrophil:lymphocyte ratio; NC = neutrophil count; LDH = lactate dehydrogenase; hs-TnI = high sensitivity troponin I; AST = aspartate aminotransferase; ALT = alanine aminotransferase; GGT = gamma-glutamyl transpeptidase; ALP = alkaline phosphatase; CKMB = creatine kinase myocardial band

Table 2: Studies that compare C-reactive protein for COVID-19

Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
Li H, et al. 2020 (39)	33.22 \pm 32.21	66.04 \pm 44.89 97.44 \pm 58.60 (critically severe)	$P = 0.001$ for all	Critically severe patients had significantly higher CRP than severe or non-severe patients. SAA correlated with CRP too consistently, indicating both should be used to reflect severity of disease – but study lacks a control group.
Liu F, et al (9)	>8.0 = 56.1% of patients 0–8.0 = 43.9% of patients	>8.0 = 93.9% of patients 0–8.0 = 6.1% of patients	$P < 0.001$ for all	Significantly more patients in the severe group experienced higher CRP levels vs non-severe.
Qin et al. 2020 (8)	33.2 (8.2–59.7)	57.9 (20.9–103.2)	$P < 0.001$	Higher levels of CRP recorded in the severe group vs non-severe group is suggestive that CRP can be monitored to assess progression of disease.
Ji et al. 2020 (10)	11.89 (9.74-23.36) (Indirect contact)	5.68 (2.80-13.0) (Direct contact)	N/A	Stratifies patients by direct and indirect contact to Wuhan – does not assess severity
Tan et al. 2020 (12)	N/A	N/A	CRP ($R = .62$; $P < .01$) ROC= 0.87 (95% CI 0.10–1.00) Cut- off= 20.42 mg/L Sensitivity=83% Specificity=91%	Absolute values not reported – instead performs ROC analysis, showing significant increase in CRP levels prior to changes in CT scores for early periods of severe group.
Wang et al. 2020	1.52 \pm 1.56 (mild)	54.15 \pm 1.06 (severe)	Mild: moderate	Groups were determined

(40)	16.76 ± 18.38 (moderate)	105.00 ± 12.73 (critical)	$P = 0.007$ Moderate: severe $P = 0.511$ Severe: critical $P = 0.947$	based on the diameter of largest lung lesion. Greater CRP values are more prominent in critical group – indicating lung damage.
------	-----------------------------	------------------------------	---	---

CRP = C-reactive protein; ROC = Receiver operating characteristic

Table 3: Studies that compare IL-6 for COVID-19

Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
Chen et al (2020) (46)	34 +/- 7	72 +/- 12	P < 0.0001	Increased expression of IL-2R and IL-6 in serum is expected to predict the severity of COVID-19
23. Liu et al. (2020) (47)	2.4 (2.1-2.9)	36.5 (30.8-42)	P < 0.0001	Severity of COVID-19 could be predicted with baseline IL-6 levels
Diao et al (2020) (48)	51 +/- 74	186 +/- 283	P < 0.0001	Significantly higher baseline levels of IL-6 in those requiring ICU compared to those who do not
Huang et al (2020) (5)	5 (0-11.2)	6.1 (1.8-37.7)	P < 0.0001	Significantly higher baseline levels of IL-6 in those requiring ICU compared to those who do not
Qin et al (2020) (8)	13.3 (3.9-41.1)	25.5 (9.5-54.5)	P < 0.0001	Significantly higher levels of IL-6 in severe and critical COVID-19. Surveillance may help in early screening of critical illness
Wu et al (2020) (49)	6.3 (5.4-7.8)	7.4 (5.6-10.9)	P < 0.0001	ARDS development in COVID-19 is related to rise in IL-6

IL-6 = interleukin-6; ARDS = Acute Respiratory Distress Syndrome; ICU = intensive care unit

Table 4: Studies that compare WCC for COVID-19

Author	Level in non-severe patient (x10 ⁹ /L)	Level in severe patient (x10 ⁹ /L)	Confidence interval (CI) range and p value	Comments
Qin et al. 2020 (8)	WCC= 4.9 (3.7–6.1) NC = 3.2 (2.1–4.4) LC = 1.0 (0.7-1.3) NLR = 3.2 (1.8–4.9)	WCC= 5.6 (4.3–8.4) NC = 4.3 (2.9–7.0) LC = 0.8 (0.6-1.1) NLR = 5.5 (3.3–10.0)	$P < 0.001$ for all	Retrospective + China + 452 Significantly higher NLR in severe patients. Monitoring may aid in early screening of critical illness.
Wang et al. 2020 (17)	WCC = 4.3 (3.3-5.4) NC = 2.7 (1.9-3.9) LC = 0.9 (0.6-1.2) (non-ICU)	WCC = 6.6 (3.6-9.8) NC = 4.6 (2.6-7.9) LC = 0.8 (0.6-1.1) (ICU)	$P = 0.003$ $P < 0.001$ $P = 0.03$	Retrospective + China + 138 Non-ICU vs ICU patients had drastically lower WCC, checking low LC and high NC may help in early detection of disease progression.
Chen N et al. 2020 (15)	WCC = 7.5 (3.5-9.5 normal range) NC = 5.0 (3.3-8.1 normal range) 38% increase LC = 0.9 (1.1-3.2 normal range) 35% decrease		N/A	Absolute values of 99 patients obtained in patients with pneumonia. Surveillance of NC/LC may reflect severity of lung abnormalities.
Tan et al. 2020 (18)	N/A		$P < 0.001$	Only lymphocyte % (LYM%) representation is graphical. TLYM % used for 90 hospitalised patients, it was able to categorise patients consistently. Surveillance may identify critically ill patients earlier.
Gong et al. 2020 (7) preprint	WBC $> 9.5 \times 10^9/L$ NC $> 7.305 \times 10^9/L$		$R = -0.54, P < 0.001$ $R = -0.585, P < 0.001$	The preprint suggests high WBC and NC indicates the higher the likelihood the severity of disease will progress to a critical stage

WCC = White cell count; NC = neutrophils; LC = lymphocytes; WBC = white blood cell

Table 5: Studies that compare LDH for COVID-19

Author	Level in non-severe patient (U/L)	Level in severe patient (U/L)	Confidence interval (CI) range and p value	Comments
Luo et al. 2020 (21)	151 (139–180)	248 (162–273)	$P < 0.002$	Higher LDH levels reported in severe patient's vs non severe.
Xiong et al. 2020 (23)	N/A	N/A	$R = 0.78$, $P < 0.001$ for correlation between LDH and CT score	LDH values linked to CT scores (used to assess severity)-no absolute values
Guan et al. 2020 (22)	≥ 250 (205/551) 37.2% of patients	≥ 250 (72/144) 58.1% of patients	N/A	Multi-centre + China + 1099 Higher LDH levels present in majority of severe patients vs non-severe. LDH measured in 675.
Mo P et al. 2020 (24)	241 (198- 338) (General)	293 (193-434) (Refractory)	$P = 0.017$	Retrospective + China + 155, refractory vs general
Ferrari D et al. 2020 (20)	276.4 ± 118.3 (COVID-19 negative)	388.0 ± 154.5 (COVID-19 positive)	$P < 0.001$	Retrospective + Italy + 141 LDH measured, COVID-19 positive vs negative
Wang et al. 2020 (17)	212 (171-291) (non-ICU)	435 (302-596) (ICU)	$P < 0.001$	Retrospective, single-centre + China + 138, non-ICU vs ICU

LDH = lactate dehydrogenase; ICU = intensive care unit

Table 6: Studies that compare D-dimer for COVID-19

Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
Tang et al. (2020) (26)	0.61 (0.35-1.29)	2.12 (0.77-5.27)	N/A	Abnormal coagulation results with markedly elevated D-dimer are common in deaths with COVID-19
Zhou et al. (2020) (32)	0.6 (0.3-1.0)	5.2 (1.5-21.1)	P < 0.001	D-dimer levels greater than 1µg/mL can help clinicians in identifying patients with poor prognosis at earlier stage
Guan et al. (2020) (22)	43.2% with > 0.5 mg/L	59.6% with > 0.5 mg/L	N/A	D-dimer levels much higher in those requiring ICU admission and invasive ventilation however statistical analysis not performed
Huang et al. (2020) (5)	0.5mg/L	2.4mg/L	P = 0.0042	Compared to non-ICU patients, ICU patients had significantly higher levels of D-dimer
Zhang et al (2020) (50)	0.41 mg/L (0.15 – 0.69)	4.76 mg/L (2.99-11.9)	P < 0.001	D-dimer on admission of greater than 2.0 µg/mL could effectively predict in-hospital mortality in patients with COVID-19 and could be an early and helpful marker to improve management

ICU = intensive care unit

Table 7: Studies that compare platelet count for COVID-19

Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
Guan et al. 2020 (22)	172,000	137,500	N/A	Although there were evident differences in lymphocytes (lymphocytopenia), platelet count (rose) and D-dimer (rose) in patients who experienced composite endpoints (ICU admission, invasive mechanical ventilation and death) there was no statistical analysis performed
Huang et al. 2020(5)	149,000	196,000	P = 0.45	Compared to non-ICU patients, ICU patients had higher plasma levels of IL-6, D-dimer and platelets. But this difference was not significant.
Liu et al. 2020 (41)	173,200 \pm 55.37	143,900 \pm 64.81	P = 0.116	C _t value of virus correlated strongly with CRP, albumin and LDH among others. Murray score for ARDS correlated with the same markers.
Liu et al. 2020 (42)	186,200	139,500	N/A	Increased CRP and decreased albumin strongly correlated with disease progression
Ruan et al. 2020 (43)	221,000 (78,000)	173,600 (67,000)	P < 0.001	Cardiac troponin, Myoglobin, CRP and IL-6 significantly increased in cases with mortality

Wang et al. 2020 (17)	165,000	142,000	$P < 0.78$	Higher WCC, specifically higher NC, lower LC and higher NLR in ICU vs non-ICU patients. Prior to the death of critical patients: NC, D-dimer, blood urea, and creatinine levels rose throughout until death, whilst the LC carried on falling. Elevated LDH also correlated with patients in ICU vs non-ICU.
Yang et al. 2020 (44)	191,000 (63,000)	164,000 (74,000)	N/A	Higher levels of platelets in severe group but lymphocytopenia seen as most common marker of infection Mortality used to indicate severity
Young et al. 2020 (45)	159,000	156,000	N/A	Lower platelets found in severe group, but statistical analysis was not performed. Severity determined by supplemental oxygen requirement
Zhou et al. 2020 (32)	220,000	165,500	$P < 0.0001$	D-dimer levels greater than $1\mu\text{g/mL}$ can help clinicians in identifying patients with poor prognosis at earlier stage There were significantly higher levels of platelets and hs-TnI in non-survivors compared to survivors Mortality used to determine

				severity
41. Liu et al. 2020 (30)	186,000 (160,000– 227,000)	105,000 (92,000– 116,000)	P < 0.001	Thrombocytopenia associated with mortality Patients split on basis of thrombocytopenia not severity
42. Yang et al. 2020 (31)	203,000 (155,000 – 257,000)	79,000 (43,000 – 129,000)	P < 0.001	Lowest platelet count associated with mortality Mortality used to indicate severity Uses nadir platelet count

IL-6 = interleukin-6; ICU = intensive care unit

Table 8: Studies that compare Cardiac Troponin Levels for COVID-19

Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
Zhou et al. (2020) (32)	3.0 (1.1-5.5)	22.2 (5.6 – 83.1)	P < 0.0001	Significantly higher levels of hs-TnI in non-survivors compared to survivors
Shi et al. (2020) (33)	<0.006 ug/L (<0.006-0.009)	0.19 ug/L (0.08-1.12)	P < 0.0001	Significantly higher levels of hs-TnI in patients who require mechanical ventilation compared to those who do not

Hs-TnI = high sensitivity troponin I

Table 9: Studies that compare creatinine levels for COVID-19


Author	Level in non-severe patient	Level in severe patient	Confidence interval (CI) range and p value	Comments
45. Xiang et al. 2020 (35)	N/A	N/A	P < 0.001	Serum values of urea, creatine and cystatin-C significantly increase in severe COVID-19 No values provided – only graphical depiction of results
46. Cheng et al. 2020 (36)	77 ± 31	132 ± 39	P < 0.001	Raised creatinine levels associated with poor outcome in COVID infection Patients split on basis of creatine level not severity
47. Zhou et al. 2020 (37)	65.3 (56.5-74.3)	71.0 (55.8-89.4)	P = 0.067	Urinalysis on admission can effectively highlight kidney impairment Patients split on basis of abnormal urinalysis

Table 10: Summary of Changes in Biomarkers Seen in Severe COVID-19 Infection

Biomarker	Change in severe COVID-19 infection
CRP	Increase
SAA	Increase
IL-6	Increase
LDH	Increase
WCC	NLR increases LC decrease
D-dimer	Increase
Platelet count	Decrease
Cardiac troponin	Increase
Renal biomarkers	Urea & creatinine increase

CRP = C-reactive protein; SAA = serum amyloid A; IL-6 = interleukin 6; LDH = lactate dehydrogenase; WCC = White cell count

Figure 1: PRISMA Flow Diagram



References

1. Gorbalenya AE, Baker SC, Baric RS, Groot RJ de, Drosten C, Gulyaeva AA, et al. Severe acute respiratory syndrome-related coronavirus: The species and its viruses – a statement of the Coronavirus Study Group. *bioRxiv*. 2020 Feb 11;2020.02.07.937862.
2. Chan JWM, Ng CK, Chan YH, Mok TYW, Lee S, Chu SYY, et al. Short term outcome and risk factors for adverse clinical outcomes in adults with severe acute respiratory syndrome (SARS). *Thorax*. 2003 Aug;58(8):686–9.
3. Booth TF, Kournikakis B, Bastien N, Ho J, Kobasa D, Stadnyk L, et al. Detection of airborne severe acute respiratory syndrome (SARS) coronavirus and environmental contamination in SARS outbreak units. *J Infect Dis*. 2005 May 1;191(9):1472–7.
4. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, Evaluation and Treatment Coronavirus (COVID-19). In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 [cited 2020 Apr 28]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK554776/>
5. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020 Feb 15;395(10223):497–506.
6. Pierce JD, McCabe S, White N, Clancy RL. Biomarkers: an important clinical assessment tool. *Am J Nurs*. 2012 Sep;112(9):52–8.

7. Gong J, Dong H, Xia SQ, Huang YZ, Wang D, Zhao Y, et al. Correlation Analysis Between Disease Severity and Inflammation-related Parameters in Patients with COVID-19 Pneumonia. medRxiv. 2020 Feb 27;2020.02.25.20025643.
8. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. Clin Infect Dis. 2020 Mar 12;
9. Liu F, Li L, Xu M, Wu J, Luo D, Zhu Y, et al. Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19. J Clin Virol. 2020 Apr 14;127:104370.
10. Ji W, Bishnu G, Cai Z, Shen X. Analysis clinical features of COVID-19 infection in secondary epidemic area and report potential biomarkers in evaluation. medRxiv. 2020 Mar 13;2020.03.10.20033613.
11. Yip TTC, Chan JWM, Cho WCS, Yip T-T, Wang Z, Kwan T-L, et al. Protein chip array profiling analysis in patients with severe acute respiratory syndrome identified serum amyloid a protein as a biomarker potentially useful in monitoring the extent of pneumonia. Clin Chem. 2005 Jan;51(1):47–55.
12. Tan C, Huang Y, Shi F, Tan K, Ma Q, Chen Y, et al. C-reactive protein correlates with computed tomographic findings and predicts severe COVID-19 early. J Med Virol. 2020 Apr 13;
13. Mahajan S, Decker CE, Yang Z, Veis D, Mellins ED, Faccio R. Plcy2/Tmem178 dependent pathway in myeloid cells modulates the pathogenesis of cytokine storm syndrome. J Autoimmun. 2019 Jun;100:62–74.

14. Mahallawi WH, Khabour OF, Zhang Q, Makhdoum HM, Suliman BA. MERS-CoV infection in humans is associated with a pro-inflammatory Th1 and Th17 cytokine profile. *Cytokine*. 2018 Apr;104:8–13.
15. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507–13.
16. Coomes EA, Haghighi H. Interleukin-6 in COVID-19: A Systematic Review and Meta-Analysis. *medRxiv*. 2020 Apr 3;2020.03.30.20048058.
17. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020 Feb 7;
18. Tan L, Wang Q, Zhang D, Ding J, Huang Q, Tang Y-Q, et al. Lymphopenia predicts disease severity of COVID-19: a descriptive and predictive study. *Signal Transduction and Targeted Therapy*. 2020 Mar 27;5(1):1–3.
19. Han Y, Zhang H, Mu S, Wei W, Jin C, Xue Y, et al. Lactate dehydrogenase, a Risk Factor of Severe COVID-19 Patients. *medRxiv*. 2020 Mar 27;2020.03.24.20040162.
20. Ferrari D, Motta A, Strollo M, Banfi G, Locatelli M. Routine blood tests as a potential diagnostic tool for COVID-19. *Clin Chem Lab Med*. 2020 Apr 16;
21. Clinical findings of 35 cases with novel coronavirus pneumonia outside of Wuhan. 2020 Apr 17 [cited 2020 Apr 29]; Available from: <https://www.researchsquare.com/article/rs-22554/v1>

22. Guan W-J, Ni Z-Y, Hu Y, Liang W-H, Ou C-Q, He J-X, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020 Feb 28;
23. Xiong Y, Sun D, Liu Y, Fan Y, Zhao L, Li X, et al. Clinical and High-Resolution CT Features of the COVID-19 Infection: Comparison of the Initial and Follow-up Changes. *Invest Radiol*. 2020 Mar 3;
24. Mo P, Xing Y, Xiao Y, Deng L, Zhao Q, Wang H, et al. Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis*. 2020 Mar 16;
25. Zhang L, Long Y, Xiao H, Yang J, Toulon P, Zhang Z. Use of D-dimer in oral anticoagulation therapy. *International Journal of Laboratory Hematology*. 2018;40(5):503–7.
26. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020;18(4):844–7.
27. Zhang L, Yan X, Fan Q, Liu H, Liu X, Liu Z, et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *J Thromb Haemost*. 2020 Apr 19;
28. Milbrandt EB, Reade MC, Lee M, Shook SL, Angus DC, Kong L, et al. Prevalence and significance of coagulation abnormalities in community-acquired pneumonia. *Mol Med*. 2009 Dec;15(11–12):438–45.

29. Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A meta-analysis. *Clin Chim Acta*. 2020 Mar 13;506:145–8.
30. Liu Y, Sun W, Guo Y, Chen L, Zhang L, Zhao S, et al. Association between platelet parameters and mortality in coronavirus disease 2019: Retrospective cohort study. *Platelets*. 2020 Apr 16;0(0):1–7.
31. Yang X, Yang Q, Wang Y, Wu Y, Xu J, Yu Y, et al. Thrombocytopenia and Its Association with Mortality in Patients with COVID-19. *Journal of Thrombosis and Haemostasis* [Internet]. [cited 2020 Apr 30];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/jth.14848>
32. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020 Mar 28;395(10229):1054–62.
33. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiol*. 2020 Mar 25;
34. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *Int Urol Nephrol* [Internet]. 2020 Mar 28 [cited 2020 Apr 30]; Available from: <https://doi.org/10.1007/s11255-020-02451-9>
35. Xiang J, Wen J, Yuan X, Xiong S, Zhou X, Liu C, et al. Potential biochemical markers to identify severe cases among COVID-19 patients. *medRxiv*. 2020 Mar 23;2020.03.19.20034447.

36. Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney International*. 2020 May 1;97(5):829–38.
37. Zhou H, Zhang Z, Fan H, Li J, Li M, Dong Y, et al. Urinalysis, but not blood biochemistry, detects the early renal-impairment in patients with COVID-19. *medRxiv*. 2020 Apr 6;2020.04.03.20051722.
38. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *Journal of Thrombosis and Haemostasis*. 2020;18(5):1094–9.
39. Li H, Xiang X, Ren H, Xu L, Zhao L, Chen X, et al. SAA is a biomarker to distinguish the severity and prognosis of Coronavirus Disease 2019 (COVID-19). *J Infect [Internet]*. 2020 Apr 8 [cited 2020 Apr 29]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7141628/>
40. Wang L. C-reactive protein levels in the early stage of COVID-19. *Med Mal Infect [Internet]*. 2020 Mar 31 [cited 2020 Apr 29]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7146693/>
41. Liu Y, Yang Y, Zhang C, Huang F, Wang F, Yuan J, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci*. 2020;63(3):364–74.
42. Liu W, Tao Z-W, Wang L, Yuan M-L, Liu K, Zhou L, et al. Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J*. 2020 May 5;133(9):1032–8.

43. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med.* 2020 Mar 3;
44. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med.* 2020 Feb 24;
45. Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA.* 2020 Mar 3;
46. Chen L, Liu HG, Liu W, Liu J, Liu K, Shang J, et al. [Analysis of clinical features of 29 patients with 2019 novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi.* 2020 Feb 6;43(0):E005.
47. Liu T, Zhang J, Yang Y, Ma H, Li Z, Zhang J, et al. The potential role of IL-6 in monitoring severe case of coronavirus disease 2019. *medRxiv.* 2020 Mar 10;2020.03.01.20029769.
48. Diao B, Wang C, Tan Y, Chen X, Liu Y, Ning L, et al. Reduction and Functional Exhaustion of T Cells in Patients with Coronavirus Disease 2019 (COVID-19). *medRxiv.* 2020 Feb 20;2020.02.18.20024364.
49. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med.* 2020 Mar 13;

50. Zhang L, Yan X, Fan Q, Liu H, Liu X, Liu Z, et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. *Journal of Thrombosis and Haemostasis* [Internet]. [cited 2020 Apr 25];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/jth.14859>

Journal Pre-proof

CRedit Author Statement

Muhammed Kermali: Writing – Original Draft, Writing – Review & Editing, Formal analysis, Visualization

Kiran Pillai: Writing – Original Draft, Writing – Review & Editing, Visualization

Raveena Kaur Khalsa: Writing – Original Draft, Writing – Review & Editing, Visualization

Zahra Ismail: Writing – Review & Editing

Amer Harky: Conceptualization, Project administration, Supervision, Visualization

Journal Pre-proof

Highlights

- There are many biomarkers with suggested roles in monitoring COVID-19 infection
- Increases in CRP and D-dimers have strong associations with mortality
- Likewise, decreases in platelet count strongly associated with mortality
- Ultimately more research is needed to determine prognostic benefit

Journal Pre-proof